

ON THE CHARACTERIZATION OF p -ADIC COLOMBEAU-EGOROV GENERALIZED FUNCTIONS BY THEIR POINT VALUES

EBERHARD MAYERHOFER

ABSTRACT. We show that contrary to recent papers by S. Albeverio, A. Yu. Khrennikov and V. Shelkovich, point values do not determine elements of the so-called p -adic Colombeau-Egorov algebra $\mathcal{G}(\mathbb{Q}_p^n)$ uniquely. We further show in a more general way that for an Egorov algebra $\mathcal{G}(M, R)$ of generalized functions on a locally compact ultrametric space (M, d) taking values in a non-trivial ring, a point value characterization holds if and only if (M, d) is discrete. Finally, following an idea due to M. Kunzinger and M. Oberguggenberger, a generalized point value characterization of $\mathcal{G}(M, R)$ is given. Elements of $\mathcal{G}(\mathbb{Q}_p^n)$ are constructed which differ from the p -adic δ -distribution considered as an element of $\mathcal{G}(\mathbb{Q}_p^n)$, yet coincide on point values with the latter.

INTRODUCTION

Algebras of generalized functions have found a growing number of applications in analysis, geometry and mathematical physics over the past two decades (cf. [3, 5, 9, 7, 6], as well as [4] for a recent unified approach). A distinguishing feature (compared to spaces of distributions in the sense of Schwartz) of Colombeau- and Egorov type algebras is the availability of a generalized point value characterization for elements of such spaces (see [10], resp. [8] for the manifold setting). Such a characterization may be viewed as a nonstandard aspect of the theory: for uniquely determining an element of a Colombeau- or Egorov algebra, its values on classical ('standard') points do not suffice: there exist elements which vanish on each classical point yet are nonzero in the quotient algebra underlying the respective construction. A unique determination can only be attained by taking into account values on generalized points, themselves given as equivalence classes of standard points. This characteristic feature is re-encountered in practically all known variants of such algebras of generalized functions.

It therefore came as a surprise when in a series of papers ([1, 2]) it was claimed that, contrary to the above general situation, in p -adic Colombeau-Egorov algebras a general point value characterization using only standard points was available. This short note is dedicated to a thorough study of (generalized) point value characterizations of p -adic Colombeau-Egorov algebras and to showing that in fact also in the p -adic setting classical point values do not suffice to uniquely determine elements of such.

1991 *Mathematics Subject Classification.* 46F30.

Key words and phrases. generalized functions, Colombeau, Egorov, generalized numbers, point values, p -adic.

Work supported by FWF research grants P16742-N04 and Y237-N13.

In the remainder of this section we recall some material from ([1, 2]), using notation from [7]. Let $\mathbb{N}$ be the natural numbers starting with $n = 1$. For a fixed prime p , let $\mathbb{Q}_p$ denote the field of rational p -adic numbers. Let $\mathcal{D}(\mathbb{Q}_p^n)$ denote the linear space of locally constant complex valued functions on $\mathbb{Q}_p^n$ ($n \geq 1$) with compact support. Let further $\mathcal{P}(\mathbb{Q}_p^n) := \mathcal{D}(\mathbb{Q}_p^n)^{\mathbb{N}}$. $\mathcal{P}(\mathbb{Q}_p^n)$ is endowed with an algebra-structure by defining addition and multiplication of sequences componentwise. Let $\mathcal{N}(\mathbb{Q}_p^n)$ be the subalgebra of elements $\{(f_k)_k\} \subset \mathcal{P}(\mathbb{Q}_p^n)$ such that for any compact set $K \subset \mathbb{Q}_p^n$ there exists an $N \in \mathbb{N}$ such that $\forall x \in K \forall k \geq N : f_k(x) = 0$. This is an ideal in $\mathcal{P}(\mathbb{Q}_p^n)$. The quotient algebra $\mathcal{G}(\mathbb{Q}_p^n) := \mathcal{P}(\mathbb{Q}_p^n)/\mathcal{N}(\mathbb{Q}_p^n)$ is called the p -adic Colombeau-Egorov algebra. Finally, so called Colombeau-Egorov generalized numbers $\tilde{\mathbb{C}}$ are introduced in the following way: Let $\tilde{\mathbb{C}}$ be the one-point compactification of $\mathbb{C} \cup \{\infty\}$.

Factorizing $\mathcal{A} = \tilde{\mathbb{C}}^{\mathbb{N}}$ by the ideal $\mathcal{I} := \{u = (u_k)_k \in \mathcal{A} \mid \exists N \in \mathbb{N} \forall k \geq N : u_k = 0\}$ yields then the ring $\tilde{\mathcal{C}}$ of Colombeau-Egorov generalized numbers. We replace $\tilde{\mathbb{C}}$ by $\mathbb{C}$ and construct similarly $\mathcal{C}$, the ring of generalized numbers: Clearly, $\tilde{\mathbb{C}}$ is not needed in this context, since representatives of elements $f \in \mathcal{G}(\mathbb{Q}_p^n)$ merely take on values in $\mathbb{C}^{\mathbb{N}}$. Let $f = [(f_k)_k] \in \mathcal{G}(\mathbb{Q}_p^n)$. It is clear that for a fixed $x \in \mathbb{Q}_p^n$, the *point value of f at x* , $[(f_k(x))_k]$ is a well defined element of $\mathcal{C}$, i.e., we may consider f as a map

$$(0.1) \quad f : \mathbb{Q}_p^n \rightarrow \mathcal{C} : x \mapsto f(x) := (f_k(x))_k + \mathcal{I}.$$

Note that the above constitutes a slight abuse of notation: The letter f denotes both a generalized function (an element of $\mathcal{G}(\mathbb{Q}_p^n)$) and a mapping on $\mathbb{Q}_p^n$.

Finally, let A be a set and let R be a ring. For $B \subset A$, $\theta \in R$ we call the characteristic function of B the map $\chi_{B,\theta} : A \rightarrow R$ which is identically θ on B and which vanishes on $A \setminus B$. Furthermore, if $\theta = 1 \in R$ we simply write $\chi_B = \chi_{B,1}$.

1. POINT VALUES AND A COUNTEREXAMPLE

The following statement is proven in Theorem 4.4 of [1]: *Let $f \in \mathcal{G}(\mathbb{Q}_p^n)$, then:*

$$f = 0 \text{ in } \mathcal{G}(\mathbb{Q}_p^n) \Leftrightarrow \forall x \in \mathbb{Q}_p^n : f(x) = 0 \text{ in } \mathcal{C}$$

However, inspired by ([11], p. 218) we construct the following counterexample to this claim, which shows that point values cannot uniquely determine elements in $\mathcal{G}(\mathbb{Q}_p^n)$ uniquely. For the sake of simplicity we assume that $n = 1$.

Example 1.1. For any $l \in \mathbb{N}$, set

$$B_l := \{x \in \mathbb{Z}_p : |x - p^l| < |p^{2l}|\} \subset \{x \in \mathbb{Z}_p : |x| = |p^l|\}.$$

For any $i \in \mathbb{N}$, we set $f_i := \chi_{B_i}$. Clearly $B_i \cap B_j = \emptyset$ whenever $i \neq j$ and since $f_i \in \mathcal{D}(\mathbb{Q}_p)$ for all natural numbers i , $(f_i)_i$ is a representative of some $f \in \mathcal{G}(\mathbb{Q}_p)$. Now, for any $\alpha \in \mathbb{Q}_p$, $f(\alpha) = 0$ in $\mathcal{C}$, since either $\alpha \in B_i$ for some $i \in \mathbb{N}$ (which implies that $f_j(\alpha) = 0 \forall j > i$) or $\alpha \in \mathbb{Q}_p \setminus \bigcup B_i$, where each f_i ($i \in \mathbb{N}$) is identically zero. Consider now the sequence $(\beta_i)_{i \geq 1} \in \mathbb{Z}_p^{\mathbb{N}}$, where $\beta_i = p^i \forall i \in \mathbb{N}$. It follows that $f_i(\beta_i) = 1 \forall i \in \mathbb{N}$. In particular, for $K = \mathbb{Z}_p$ or any clopen ball containing 0, there is no representative $(g_j)_j$ of f such that for some $N > 0$, $g_j = 0 \forall j \geq N$. Hence $f \neq 0$ in $\mathcal{G}(\mathbb{Q}_p)$ although all standard point values of f vanish.

Remark 1.2. *By means of the above example we may analyze the proof of Theorem 4.4 in [1]. Let f be the generalized function from 1.1. As a compact set choose $K :=$*

$B_{\leq p^{-2}}(0) = p^2\mathbb{Z}_p$. For the representative $(f_k)_k$ constructed in 1.1 and $x = 0$ we have $N(0) = 1$, which in the notation of [1] means that for any $k \geq 1 = N(0)$, $f_k(0) = 0$. Also, recall that $B_\gamma(a)$ is the dressed ball $B_{\leq p^\gamma}(a)$. The “parameter of constancy” ([1], p. 6) of f_1 at $x = 0$, which is the maximal γ such that f_1 is identically zero on $B_\gamma(0)$, is $l_0(0) = -2$. Now, there exists a covering of K consisting of a single set, namely $B_{l_0(0)}(0)$. Thus we may replace the application of the Heine-Borel Lemma in [1] by our singleton-covering. But then the claim that (4.1) and (4.2) imply that for all $k \geq N(0) = 1$ we have $f_k(0 + x') = f_k(0) = 0 \forall x' \in K$ does not hold. This indeed follows from the definition of the sequence $(f_k)_k$ of locally constant functions from above, since for any $k \in \mathbb{N}$ we have $f_k(p^k) = 1$.

2. EGOROV ALGEBRAS ON LOCALLY COMPACT ULTRAMETRIC SPACES

In this section we consider the problem of point value characterization in Egorov algebras in full generality: to this end we consider a general locally compact ultrametric space (M, d) instead of $\mathbb{Q}_p^n$, where M need not have a field structure. Our aim is to show that even in such a general setting, the respective algebra cannot have a point value characterization, unless M carries the discrete topology. Denote by $\mathcal{E}_d(M)$ the algebra of sequences of locally constant functions with compact support, taking values in a commutative ring $R \neq \{0\}$. Let $\mathcal{N}_d(M)$ be the set of negligible functions $\{(f_k)_k\} \subset \mathcal{E}_d(M)$ such that for any compact set $K \subset M$ there exists an $N \in \mathbb{N}$ such that $\forall x \in K \forall k \geq N : f_k(x) = 0$. The subset $\mathcal{N}_d(M)$ is an ideal in $\mathcal{E}_d(M)$ and the quotient algebra $\mathcal{G}(M, R) := \mathcal{E}_d(M)/\mathcal{N}_d(M)$ is called the ultrametric Egorov algebra associated with (M, d) . Furthermore, the ring of generalized numbers is defined by $\mathcal{R} := \mathbb{R}^\mathbb{N}/\sim$, where $\sim$ is the equivalence relation on $\mathbb{R}^\mathbb{N}$ given by

$$u \sim v \text{ in } \mathbb{R}^\mathbb{N} \Leftrightarrow \exists N \in \mathbb{N} \forall k \geq N : u_k - v_k = 0.$$

We call $\mathcal{I}(R) := \{w \in R^\mathbb{N} : w \sim 0\}$ the ideal of negligible sequences in R . Analogous to (0.1), for $f \in \mathcal{G}(M, R)$ evaluation on standard point values is introduced by means of the mapping:

$$(2.2) \quad f : M \rightarrow \mathcal{R} : x \mapsto f(x) := (f_k(x))_k + \mathcal{I}(R).$$

Definition 2.1. An ultrametric Egorov algebra $\mathcal{G}(M, R)$ is said to admit a standard point value characterization if for each $u \in \mathcal{G}(M, R)$ we have

$$u = 0 \Leftrightarrow \forall x \in M : u(x) = 0 \text{ in } \mathcal{R}$$

Using this terminology, Example 1.1 shows that $\mathcal{G}(\mathbb{Q}_p^n)$ does not admit a standard point value characterization. The main result of this section is:

Theorem 2.2. Let (M, d) be a locally compact ultrametric space and let $R \neq \{0\}$. Then $\mathcal{G}(M, R)$ does not admit a standard point value characterization unless (M, d) is discrete.

Proof. The result follows by generalizing the construction of Example 1.1. Assume (M, d) is not discrete, then there exists a point $x \in M$ and a sequence $(x_n)_n$ of distinct points in M converging to x . We may assume that $d(x, x_i) > d(x, x_j)$ whenever $i < j$. Define stripped balls $(B_n)_{n \geq 1}$ with centers $(x_n)_{n \geq 1}$ by $B_n := \{y \in M \mid d(x_n, y) < \frac{d(x_n, x)}{2}\}$. Due to the ultrametric property “the strongest one wins” we have $B_n \subset \{z \mid d(x, z) = d(x_n, x)\}$, which further implies that for all $i \neq j$, $i, j \in \mathbb{N}$ the balls B_i, B_j are disjoint sets in M . Since R is a non-trivial

ring, we may choose some $\theta \in R \setminus \{0\}$. Now we define a sequence $(f_k)_k$ of locally constant functions in the following way: For any $i \geq 1$ set $f_i = \chi_{B_i, \theta}$. Clearly, $f := [(f_i)_i] \in \mathcal{G}(M, R)$, and similarly to Example 1.1, for any $\alpha \in M$, $f(\alpha) = 0$ in $\mathcal{R}$. Nevertheless for the sequence $(x_n)_n$, which without loss of generality may be assumed to lie in a compact neighborhood of x , one has $f_i(x_i) = \theta \forall i \geq 1$ which implies that $f \neq 0$ in $\mathcal{G}(M, R)$. $\square$

Recall that a discrete topological space X has the following properties:

- (i) X is locally compact.
- (ii) Any compact set in X contains finitely many points only.

Therefore we know that for a set D endowed with the discrete metric and for any commutative ring R , the respective ultrametric Egorov algebra $\mathcal{G}(D, R)$ admits a pointwise characterization. We therefore conclude:

Corollary 2.3. *For a locally compact ultrametric space (M, d) and a non-trivial ring R , the following statements are equivalent:*

- (i) $\mathcal{G}(M, R)$ admits a standard point value characterization.
- (ii) The topology of (M, d) is discrete.

3. GENERALIZED POINT VALUES

In this section we give an appropriate generalized point value characterization in the style of ([7], pp. 37–43) of $\mathcal{G}(M, R)$, where M is endowed with a non-discrete ultrametric d for which M is locally compact, and $R \neq \{0\}$. First, we have to introduce a set $\widetilde{M}_c$ of compactly supported generalized points over M . Let $\mathcal{E} = M^{\mathbb{N}}$, the ring of sequences in M , and identify two sequences, if for some index $N \in \mathbb{N}$ one has $d(x_n, y_n) = 0 \forall n \geq N$, i.e., $x_n = y_n \forall n \geq N$; we write $x \sim y$. We call $\widetilde{M} = \mathcal{E} / \sim$ the ring of generalized numbers. Finally, $\widetilde{M}_c$ is the subset of such elements $x \in \widetilde{M}$ for which there exists a compact subset K and some representative $(x_n)_n$ of x such that for some $N > 0$ we have $x_n \in K$ for all $n \geq N$. It follows that evaluating a function $u \in \mathcal{G}(M, R)$ at a compactly supported generalized point x is possible, i.e., for representatives $(x_k)_k$, $(u_k)_k$ of x resp. u , $[(u_k(x_k))_k]$ is a well defined element of $\mathcal{R}$.

Proposition 3.1. *In $\mathcal{G}(M, R)$, there is a generalized point value characterization, i.e.,*

$$u = 0 \text{ in } \mathcal{G}(M, R) \Leftrightarrow \forall x \in \widetilde{M}_c : u(x) = 0 \text{ in } \mathcal{R}$$

Proof. The condition is obviously necessary. Conversely, let $u \in \mathcal{G}(M, R)$, $u \neq 0$. This means that there is a representative $(u_k)_k$ of u and a compact set $K \subset\subset M$ such that u_k does not vanish on K for infinitely many $k \in \mathbb{N}$. In particular this means we have a sequence $(x_k)_k$ in K such that for infinitely many $k \in \mathbb{N}$, $u_k(x_k) \neq 0$. Clearly this means that $u(x) \neq 0$ in $\mathcal{R}$ for the compactly supported generalized point defined by $x := [(x_k)_k]$. $\square$

4. THE IMBEDDED δ -DISTRIBUTION IN $\mathcal{G}(\mathbb{Q}_p^n)$.

In [1], Theorem 4.4 is illustrated by some examples, to highlight the advantage of a point value concept in $\mathcal{G}(\mathbb{Q}_p^n)$. In this section we discuss the δ -distribution (Example 4.5 on p. 12 in [1]) and construct a generalized function $f \in \mathcal{G}(\mathbb{Q}_p)$ different from δ which however coincides with δ on all standard points in $\mathbb{Q}_p$. We

first imbed the δ -distribution in $\mathcal{G}(\mathbb{Q}_p)$ as in [1] (p. 9, Theorem 3.3) which yields $\iota(\delta) = (\delta_k)_k + \mathcal{N}_p(\mathbb{Q}_p)$, where $\delta_k(x) := p^k \Omega(p^k |x|_p)$ and Ω is the bump function on $\mathbb{R}_0^+$ given by

$$\Omega(t) := \begin{cases} 1, & 0 \leq t \leq 1 \\ 0, & t > 1 \end{cases}$$

Evaluation of $\iota(\delta)$ on standard points is shown in Example 4.5 of [1]. With $\tilde{c} := (p^k)_k + \mathcal{I} \in \mathcal{C}$ one has:

$$\iota(\delta)(x) = \begin{cases} \tilde{c}, & x = 0 \\ 0, & x \neq 0 \end{cases} \quad (x \in \mathbb{Q}_p)$$

Let $\varphi : \mathbb{N} \rightarrow \mathbb{Z}$ be a monotonous function such that $\lim_{k \rightarrow \infty} \varphi(k) = \infty$, and such that the cardinality of $U_\varphi := \{k : \varphi(k) > k\}$ is infinite. Consider an element $f \in \mathcal{G}(\mathbb{Q}_p)$ given by $f := (f_k)_k + \mathcal{N}(\mathbb{Q}_p)$ where for any $k \geq 1$, $f_k(x) := p^k \Omega(p^{\varphi(k)} |x|_p)$. Then the standard point values of $\iota(\delta)$ and f coincide. Furthermore, they coincide on compactly supported generalized points $x \in \tilde{\mathbb{Q}}_{p,c}$ with the property that for any representative $(x_k)_k$ of x there exists an $N \in \mathbb{N}$ such that $\forall k \geq N : |x_k|_p > p^{-\min\{k, \varphi(k)\}}$, since in this case we have $\delta_k(x_k) = f_k(x_k) = 0$. However, there are compactly supported generalized points violating this condition which yield different generalized point values of $\iota(\delta)$ resp. f : consider the generalized point $x_0 := [(p^k)_k] \in \tilde{\mathbb{Q}}_{p,c}$. Then $f(x_0) \neq \tilde{c}$, since $\theta_k := f_k(x_k) = 0$ for any $k \in U_\varphi$ and thus $\theta_k = 0$ for infinitely many $k \in \mathbb{N}$. But $\iota(\delta)(x_0) = \tilde{c}$.

REFERENCES

- [1] S. ALBEVERIO, A. Y. KHRENNIKOV, AND V. M. SHELKOVICH, *Nonlinear singular problems of p-adic analysis: associative algebras of p-adic distributions*, Izv. Ross. Akad. Nauk Ser. Mat., 69 (2005), pp. 3–44.
- [2] S. ALBEVERIO, A. Y. KHRENNIKOV, AND V. M. SHELKOVICH, *p-adic Colombeau-Egorov type theory of generalized functions*, Math. Nachr., 278 (2005), pp. 3–16.
- [3] J. F. COLOMBEAU *Elementary introduction to new generalized functions*. North-Holland, Amsterdam, 1985.
- [4] A. DELCROIX, M. HASLER, S. PILIPOVIĆ, V. VALMORIN, *Generalized function algebras as sequence space algebras*, Proc. Amer. Math. Soc. 132 (2004), no. 7, 2031–2038.
- [5] YU. V. EGOROV, *A contribution to the theory of generalized functions*, Russ. Math. Surveys 45:5 (1990), 1–49.
- [6] M. KUNZINGER, *Nonsmooth differential geometry and algebras of generalized functions*, J. Math. Anal. Appl. 297 (2004), no. 2, 456–471.
- [7] M. GROSSER, M. KUNZINGER, M. OBERGUGGENBERGER, AND R. STEINBAUER, *Geometric theory of generalized functions with applications to general relativity*, vol. 537 of Mathematics and its Applications, Kluwer Academic Publishers, Dordrecht, 2001.
- [8] M. KUNZINGER, R. STEINBAUER, *Foundations of a nonlinear distributional geometry*, Acta Appl. Math. 71 (2002), no. 2, 179–206.
- [9] M. OBERGUGGENBERGER *Multiplication of distributions and applications to partial differential equations*. Pitman Research Notes in Mathematics 259, Longman, Harlow, 1992.
- [10] M. OBERGUGGENBERGER AND M. KUNZINGER, *Characterization of Colombeau generalized functions by their pointvalues*, Math. Nachr., 203 (1999), 147–157.
- [11] A. M. ROBERT, *A course in p-adic analysis*, vol. 198 of Graduate Texts in Mathematics, Springer-Verlag, New York, 2000.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIENNA, NORDBERGSTRASSE 15, 1090 VIENNA, AUSTRIA

E-mail address: eberhard.mayerhofer@univie.ac.at